

# On Ostrowski type inequalities for Stieltjes integrals with absolutely continuous integrands and integrators of bounded variation

P. Cerone<sup>a,\*</sup>, W.S. Cheung<sup>b</sup>, S.S. Dragomir<sup>a</sup>

<sup>a</sup>*School of Computer Science and Mathematics, Victoria University, PO Box 14428, Melbourne City, VIC 8001, Australia*

<sup>b</sup>*Department of Mathematics, University of Hong Kong, Pokfulam Road, Hong Kong, China*

Received 13 October 2006; received in revised form 7 December 2006; accepted 12 December 2006

## Abstract

Some Ostrowski type inequalities are given for the Stieltjes integral where the integrand is absolutely continuous while the integrator is of bounded variation. The case when  $|f'|$  is convex is explored. Applications for the mid-point rule and a generalised trapezoid type rule are also presented.

© 2007 Elsevier Ltd. All rights reserved.

**Keywords:** Integral inequalities; Ostrowski inequality; Stieltjes integral; Quadrature rules; Mid-point rule; Trapezoidal rule; Convex functions

## 1. Introduction

The following result is known in the literature as *Ostrowski's inequality* [1]:

Let  $f : [a, b] \rightarrow \mathbb{R}$  be a differentiable mapping on  $(a, b)$  with the property that  $|f'(t)| \leq M$  for all  $t \in (a, b)$ . Then

$$\left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \left[ \frac{1}{4} + \left( \frac{x - \frac{a+b}{2}}{b-a} \right)^2 \right] (b-a) M \quad (1.1)$$

for all  $x \in (a, b)$ . The constant  $\frac{1}{4}$  is best possible in the sense that it cannot be replaced by a smaller constant.

The above result has been naturally extended for absolutely continuous functions and Lebesgue  $p$ -norms of the derivative  $f'$  in [2–4]. They can also be obtained, in a slightly different form, as particular cases of some results established by Fink in [5] for  $n$ -time differentiable functions.

For other Ostrowski type inequalities concerning Lipschitzian and  $r - H$ -Hölder type functions, see [6,7].

\* Corresponding author.

E-mail addresses: [pietro.cerone@vu.edu.au](mailto:pietro.cerone@vu.edu.au) (P. Cerone), [wscheung@hkucc.hku.hk](mailto:wscheung@hkucc.hku.hk) (W.S. Cheung), [sever.dragomir@vu.edu.au](mailto:sever.dragomir@vu.edu.au) (S.S. Dragomir).

URLs: <http://rgmia.vu.edu.au/cerone> (P. Cerone), <http://rgmia.vu.edu.au/dragomir> (S.S. Dragomir).

The cases of bounded variation functions and monotonic functions were considered in [8,9] while the case of convex functions was studied in [10].

In an effort to obtain an Ostrowski type inequality for the Stieltjes integral, which obviously contains the weighted integrals case, Dragomir established in [11] the following result:

**Theorem 1.** Let  $f : [a, b] \rightarrow \mathbb{R}$  be a function of bounded variation and  $u : [a, b] \rightarrow \mathbb{R}$  a function of  $r$  –  $H$ -Hölder type, i.e.,

$$|u(x) - u(y)| \leq H|x - y|^r \quad \text{for any } x, y \in [a, b], \quad (1.2)$$

where  $r \in (0, 1]$  and  $H > 0$  are given. Then, for any  $x \in [a, b]$ ,

$$\begin{aligned} \left| [u(b) - u(a)] f(x) - \int_a^b f(t) du(t) \right| &\leq H \left[ (x-a)^r \bigvee_a^x(f) + (b-x)^r \bigvee_x^b(f) \right] \\ &\leq H \times \begin{cases} [(x-a)^r + (b-x)^r] \left[ \frac{1}{2} \bigvee_a^b(f) + \frac{1}{2} \left| \bigvee_a^x(f) - \bigvee_x^b(f) \right| \right]; \\ [(x-a)^{qr} + (b-x)^{qr}]^{\frac{1}{q}} \left[ \left( \bigvee_a^x(f) \right)^p + \left( \bigvee_x^b(f) \right)^p \right]^{\frac{1}{p}} & \text{if } p > 1, \frac{1}{p} + \frac{1}{q} = 1; \\ \left[ \frac{1}{2}(b-a) + \left| x - \frac{a+b}{2} \right| \right]^r \bigvee_a^b(f), \end{cases} \end{aligned} \quad (1.3)$$

where  $\bigvee_c^d(f)$  denotes the total variation of  $f$  on the interval  $[c, d]$ .

The dual case was considered in [12] and can be stated as follows:

**Theorem 2.** Let  $u : [a, b] \rightarrow \mathbb{R}$  be a function of bounded variation on  $[a, b]$  and  $f : [a, b] \rightarrow \mathbb{R}$  a function of  $r$  –  $H$ -Hölder type. Then

$$\left| [u(b) - u(a)] f(x) - \int_a^b f(t) du(t) \right| \leq H \left[ \frac{1}{2}(b-a) + \left| x - \frac{a+b}{2} \right| \right]^r \bigvee_a^b(u) \quad (1.4)$$

for any  $x \in [a, b]$ .

For other results concerning inequalities for Stieltjes integrals, see [13–16].

The aim of the present paper is to continue the study of Ostrowski type inequalities for Stieltjes integrals  $\int_a^b f(t) du(t)$  where the function  $f$ , the *integrand*, is assumed to be absolutely continuous while the *integrator*  $u$ , is of bounded variation. Applications to the mid-point rule and for a generalised trapezoid rule are also pointed out.

## 2. General bounds for absolutely continuous functions

The following representation result is of interest:

**Lemma 1.** Let  $f : [a, b] \rightarrow \mathbb{R}$  be an absolutely continuous function on  $[a, b]$  and  $u : [a, b] \rightarrow \mathbb{R}$  such that the Stieltjes integral  $\int_a^b f(t) du(t)$  exists. Then

$$f(x)[u(b) - u(a)] - \int_a^b f(t) du(t) = \int_a^b (x-t) \left( \int_0^1 f'[\lambda t + (1-\lambda)x] d\lambda \right) du(t) \quad (2.1)$$

or, equivalently,

$$\begin{aligned} \int_a^b u(t) df(t) - u(b)[f(b) - f(x)] - u(a)[f(x) - f(a)] \\ = \int_a^b (x-t) \left( \int_0^1 f'[\lambda t + (1-\lambda)x] d\lambda \right) du(t) \end{aligned} \quad (2.2)$$

for each  $x \in [a, b]$ .

**Proof.** Since  $f$  is absolutely continuous on  $[a, b]$ , hence, for any  $x, t \in [a, b]$  with  $x \neq t$ , one has

$$\frac{f(x) - f(t)}{x - t} = \frac{\int_t^x f'(u) du}{x - t} = \int_0^1 f'[(1 - \lambda)x + \lambda t] d\lambda$$

giving the equality of interest (see also [17]):

$$f(x) = f(t) + (x - t) \int_0^1 f'[(1 - \lambda)x + \lambda t] d\lambda \quad (2.3)$$

for any  $x, t \in [a, b]$ .

Integrating the identity (2.3) we deduce

$$f(x) \int_a^b du(t) = \int_a^b f(t) du(t) + \int_a^b (x - t) \left( \int_0^1 f'[(1 - \lambda)x + \lambda t] d\lambda \right) du(t),$$

which is exactly the desired representation (2.1).

Now, on utilising the integration by parts formula for the Stieltjes integral in the left side of (2.1), we obtain (2.2). ■

For an absolutely continuous function  $f : [a, b] \rightarrow \mathbb{R}$ , let us denote by  $\mu(f; x, t) := \left| \int_0^1 f'[\lambda t + (1 - \lambda)x] d\lambda \right|$ , where  $(t, x) \in [a, b]^2$ . It is obvious that, by the Hölder inequality, we have

$$\mu(f; x, t) \leq \begin{cases} \|f'\|_{[t, x], \infty} & \text{if } f' \in L_\infty[a, b]; \\ \|f'\|_{[t, x], p} & \text{if } f' \in L_p[a, b], p \geq 1, \end{cases} \quad (2.4)$$

where

$$\|f'\|_{[t, x], \infty} := \sup_{\substack{u \in [t, x] \\ (u \in [x, t])}} |f'(u)|,$$

$$\|f'\|_{[t, x], p} := \left| \int_t^x |f'(u)|^p du \right|^{\frac{1}{p}}, \quad p \geq 1$$

and  $t, x \in [a, b]$ .

We can state now the following inequality of Ostrowski type for the Stieltjes integral that provides various bounds which complement the results in Theorems 1 and 2 from the introduction:

**Theorem 3.** Let  $f : [a, b] \rightarrow \mathbb{R}$  be an absolutely continuous function and  $u : [a, b] \rightarrow \mathbb{R}$  a function of bounded variation on  $[a, b]$ . Then

$$\left| [u(b) - u(a)] f(x) - \int_a^b f(t) du(t) \right| \leq M(x), \quad (2.5)$$

and, equivalently,

$$\left| \int_a^b u(t) df(t) - u(b)[f(b) - f(x)] - u(a)[f(x) - f(a)] \right| \leq M(x), \quad (2.6)$$

where  $M(x) = M_1(x) + M_2(x)$  and

$$M_1(x) := \bigvee_a^x(u) \sup_{t \in [a, x]} [(x - t) \mu(f; x, t)],$$

$$M_2(x) := \bigvee_x^b(u) \sup_{t \in [x, b]} [(t - x) \mu(f; x, t)],$$

for  $x \in [a, b]$ .

**Proof.** We use the fact that, if  $p, v : [c, d] \rightarrow \mathbb{R}$  are such that  $p$  is continuous and  $v$  is of bounded variation, then the Stieltjes integral  $\int_c^d p(t) dv(t)$  exists and

$$\left| \int_c^d p(x) dv(x) \right| \leq \sup_{x \in [c, d]} |p(x)| \bigvee_c^d(v).$$

Utilising the representation (2.1) we have

$$\begin{aligned} \left| f(x)[u(b) - u(a)] - \int_a^b f(t) du(t) \right| &= \left| \int_a^x (x-t) \left( \int_0^1 f'[\lambda t + (1-\lambda)x] d\lambda \right) du(t) \right. \\ &\quad \left. + \int_x^b (x-t) \left( \int_0^1 f'[\lambda t + (1-\lambda)x] d\lambda \right) du(t) \right| \\ &\leq \left| \int_a^x (x-t) \left( \int_0^1 f'[\lambda t + (1-\lambda)x] d\lambda \right) du(t) \right| \\ &\quad + \left| \int_x^b (x-t) \left( \int_0^1 f'[\lambda t + (1-\lambda)x] d\lambda \right) du(t) \right| \\ &\leq \bigvee_a^x(u) \sup_{t \in [a, x]} [(x-t) \mu(f; x, t)] \\ &\quad + \bigvee_x^b(u) \sup_{t \in [x, b]} [(t-x) \mu(f; x, t)], \end{aligned}$$

and the theorem is proved. ■

**Remark 1.** Using the notations in Theorem 3, we have

$$\begin{aligned} M_1(x) &\leq (x-a) \bigvee_a^x(u) \sup_{t \in [a, x]} \mu(f; x, t) \\ &\leq (x-a) \bigvee_a^x(u) \cdot \begin{cases} \|f'\|_{[a, x], \infty} & \text{if } f' \in L_\infty[a, b]; \\ \|f'\|_{[a, x], p} & \text{if } f' \in L_p[a, b], p \geq 1, \end{cases} \end{aligned}$$

and

$$\begin{aligned} M_2(x) &\leq (b-x) \bigvee_x^b(u) \sup_{t \in [x, b]} \mu(f; x, t) \\ &\leq (b-x) \bigvee_x^b(u) \cdot \begin{cases} \|f'\|_{[x, b], \infty} & \text{if } f' \in L_\infty[a, b]; \\ \|f'\|_{[x, b], p} & \text{if } f' \in L_p[a, b], p \geq 1, \end{cases} \end{aligned}$$

for any  $x \in [a, b]$ .

The above inequalities for  $M_1$  and  $M_2$  are obvious from the inequality (2.4) and the details are omitted.

**Remark 2.** If we denote by  $\|f'\|_{[c, d], p}$  the  $p$  norm on the interval  $[c, d]$ , where  $1 \leq p \leq \infty$ , then for  $f' \in L_p[a, b]$ , we have

$$\left| f(x)[u(b) - u(a)] - \int_a^b f(t) du(t) \right| \leq (x-a) \bigvee_a^x(u) \|f'\|_{[a, x], p} + (b-x) \bigvee_x^b(u) \|f'\|_{[x, b], p} =: N(x), \quad (2.7)$$

where  $p \in [1, \infty]$  and  $x \in [a, b]$ .

Obviously one can derive many upper bounds for the function  $N(x)$  defined above. We intend to present in the following only a few that are simple and perhaps of interest for applications.

**Estimate 1.** We have

$$\begin{aligned}
 N(x) &\leq \left[ (x-a) \bigvee_a^x(u) + (b-x) \bigvee_x^b(u) \right] \|f'\|_{[a,b],p} \\
 &\leq \|f'\|_{[a,b],p} \cdot \begin{cases} \left[ \frac{1}{2}(b-a) + \left| x - \frac{a+b}{2} \right| \right] \bigvee_a^b(u); \\ \left[ (x-a)^\alpha + (b-x)^\alpha \right]^{\frac{1}{\alpha}} \left[ \left( \bigvee_a^x(u) \right)^\beta + \left( \bigvee_x^b(u) \right)^\beta \right]^{\frac{1}{\beta}} & \text{if } \alpha > 1, \frac{1}{\alpha} + \frac{1}{\beta} = 1; \\ (b-a) \left[ \frac{1}{2} \bigvee_a^b(u) + \frac{1}{2} \left| \bigvee_a^x(u) - \bigvee_x^b(u) \right| \right] \end{cases} \quad (2.8)
 \end{aligned}$$

for any  $x \in [a, b]$ .

**Estimate 2.** We have

$$\begin{aligned}
 N(x) &\leq \left[ \frac{1}{2}(b-a) + \left| x - \frac{a+b}{2} \right| \right] \left[ \bigvee_a^x(u) \|f'\|_{[a,x],p} + \bigvee_x^b(u) \|f'\|_{[x,b],p} \right] \\
 &\leq \left[ \frac{1}{2}(b-a) + \left| x - \frac{a+b}{2} \right| \right] \\
 &\quad \times \begin{cases} \max \{ \|f'\|_{[a,x],p}, \|f'\|_{[x,b],p} \} \bigvee_a^b(u); \\ \|f'\|_{[a,b],p} \left[ \left( \bigvee_a^x(u) \right)^q + \left( \bigvee_x^b(u) \right)^q \right]^{\frac{1}{q}} & \text{if } p > 1, \frac{1}{p} + \frac{1}{q} = 1; \\ \left[ \frac{1}{2} \bigvee_a^b(u) + \frac{1}{2} \left| \bigvee_a^x(u) - \bigvee_x^b(u) \right| \right] [\|f'\|_{[a,x],p} + \|f'\|_{[x,b],p}] \end{cases}
 \end{aligned}$$

for any  $x \in [a, b]$ .

**Estimate 3.** We have

$$\begin{aligned}
 N(x) &\leq \left[ \frac{1}{2} \bigvee_a^b(u) + \frac{1}{2} \left| \bigvee_a^x(u) - \bigvee_x^b(u) \right| \right] \left[ (x-a) \|f'\|_{[a,x],p} + (b-x) \|f'\|_{[x,b],p} \right] \\
 &\leq \left[ \frac{1}{2} \bigvee_a^b(u) + \frac{1}{2} \left| \bigvee_a^x(u) - \bigvee_x^b(u) \right| \right] \\
 &\quad \times \begin{cases} \max \{ \|f'\|_{[a,x],p}, \|f'\|_{[x,b],p} \} (b-a); \\ \left[ (x-a)^q + (b-x)^q \right]^{\frac{1}{q}} \|f'\|_{[a,b],p} & \text{if } p > 1, \frac{1}{p} + \frac{1}{q} = 1; \\ \left[ \frac{1}{2}(b-a) + \left| x - \frac{a+b}{2} \right| \right] [\|f'\|_{[a,x],p} + \|f'\|_{[x,b],p}] \end{cases}
 \end{aligned}$$

for each  $x \in [a, b]$ .

In practical applications, the mid-point rule, that results for  $x = \frac{a+b}{2}$ , is of obvious interest due to its simpler form.

**Corollary 1.** With the assumptions in Theorem 3, we have the inequalities:

$$\left| [u(b) - u(a)] f\left(\frac{a+b}{2}\right) - \int_a^b f(t) \, du(t) \right| \leq \frac{1}{2}(b-a) \left[ \bigvee_a^{\frac{a+b}{2}}(u) \|f'\|_{[a, \frac{a+b}{2}],p} + \bigvee_{\frac{a+b}{2}}^b(u) \|f'\|_{[\frac{a+b}{2}, b],p} \right]$$

$$\leq \frac{1}{2} (b-a) \begin{cases} \max \left\{ \|f'\|_{[a, \frac{a+b}{2}], p}, \|f'\|_{[\frac{a+b}{2}, b], p} \right\} \bigvee_a^b(u); \\ \left[ \|f'\|_{[a, \frac{a+b}{2}], p}^\alpha + \|f'\|_{[\frac{a+b}{2}, b], p}^\alpha \right]^{\frac{1}{\alpha}} \left[ \left( \bigvee_a^{\frac{a+b}{2}}(u) \right)^\beta + \left( \bigvee_{\frac{a+b}{2}}^b(u) \right)^\beta \right]^{\frac{1}{\beta}} \\ \text{if } \alpha > 1, \frac{1}{\alpha} + \frac{1}{\beta} = 1; \\ \left[ \frac{1}{2} \bigvee_a^b(u) + \frac{1}{2} \left| \bigvee_a^{\frac{a+b}{2}}(u) - \bigvee_{\frac{a+b}{2}}^b(u) \right| \right] \left[ \|f'\|_{[a, \frac{a+b}{2}], p} + \|f'\|_{[\frac{a+b}{2}, b], p} \right], \end{cases} \quad (2.9)$$

where  $p \in [1, \infty]$ .

From the above, it is obvious that we can get some appealing inequalities as follows:

$$\begin{aligned} & \left| [u(b) - u(a)] f\left(\frac{a+b}{2}\right) - \int_a^b f(t) du(t) \right| \\ & \leq \frac{1}{2} (b-a) \begin{cases} \|f'\|_{[a,b], \infty} \bigvee_a^b(u), & \text{if } f' \in L_\infty[a, b]; \\ \|f'\|_{[a,b], p} \left[ \left( \bigvee_a^{\frac{a+b}{2}}(u) \right)^q + \left( \bigvee_{\frac{a+b}{2}}^b(u) \right)^q \right]^{\frac{1}{q}} & \text{if } p > 1, \frac{1}{p} + \frac{1}{q} = 1, f' \in L_p[a, b]; \\ \left[ \frac{1}{2} \bigvee_a^b(u) + \frac{1}{2} \left| \bigvee_a^{\frac{a+b}{2}}(u) - \bigvee_{\frac{a+b}{2}}^b(u) \right| \right] \|f'\|_{[a,b], 1}. \end{cases} \quad (2.10) \end{aligned}$$

**Remark 3.** Similar inequalities can be obtained for the generalised trapezoid rule. We only state here the following simple result:

$$\begin{aligned} & \left| \int_a^b u(t) df(t) - u(b) \left[ f(b) - f\left(\frac{a+b}{2}\right) \right] - u(a) \left[ f\left(\frac{a+b}{2}\right) - f(a) \right] \right| \\ & \leq \frac{1}{2} (b-a) \begin{cases} \|f'\|_{[a,b], \infty} \bigvee_a^b(u), & \text{if } f' \in L_\infty[a, b]; \\ \|f'\|_{[a,b], p} \left[ \left( \bigvee_a^{\frac{a+b}{2}}(u) \right)^q + \left( \bigvee_{\frac{a+b}{2}}^b(u) \right)^q \right]^{\frac{1}{q}} & \text{if } p > 1, \frac{1}{p} + \frac{1}{q} = 1, f' \in L_p[a, b]; \\ \left[ \frac{1}{2} \bigvee_a^b(u) + \frac{1}{2} \left| \bigvee_a^{\frac{a+b}{2}}(u) - \bigvee_{\frac{a+b}{2}}^b(u) \right| \right] \|f'\|_{[a,b], 1} \end{cases} \end{aligned}$$

provided that  $u$  is of bounded variation and  $f$  is absolutely continuous on  $[a, b]$ .

### 3. Bounds in the case of $|f'|$ a convex function

Some of the above results can be improved provided that a convexity assumption for  $|f'|$  is in place:

**Theorem 4.** Let  $f : [a, b] \rightarrow \mathbb{R}$  be an absolutely continuous function on  $[a, b]$ ,  $u : [a, b] \rightarrow \mathbb{R}$  a function of bounded variation on  $[a, b]$  and  $x \in [a, b]$ . If  $|f'|$  is convex on  $[a, x]$  and  $[x, b]$  (and the intervals can be reduced to a single

point), then

$$\begin{aligned}
 & \left| [u(b) - u(a)] f(x) - \int_a^b f(t) du(t) \right| \\
 & \leq \frac{1}{2} \left[ \bigvee_a^x(u) \sup_{t \in [a, x]} \{ (x-t) |f'(t)| \} + \bigvee_x^b(u) \sup_{t \in [x, b]} \{ (t-x) |f'(t)| \} \right] \\
 & \quad + \frac{1}{2} |f'(x)| \left[ (x-a) \bigvee_a^x(u) + (b-x) \bigvee_x^b(u) \right] \\
 & \leq \frac{1}{2} \left[ (x-a) \bigvee_a^x(u) \|f'\|_{[a, x], \infty} + (b-x) \bigvee_x^b(u) \|f'\|_{[x, b], \infty} \right] \\
 & \quad + \frac{1}{2} |f'(x)| \left[ (x-a) \bigvee_a^x(u) + (b-x) \bigvee_x^b(u) \right], \tag{3.1}
 \end{aligned}$$

for any  $x \in [a, b]$ .

**Proof.** As in the proof of Theorem 3, we have

$$\begin{aligned}
 & \left| f(x) [u(b) - u(a)] - \int_a^b f(t) du(t) \right| \leq \sup_{t \in [a, x]} \left[ (x-t) \left| \int_0^1 f'[\lambda t + (1-\lambda)x] d\lambda \right| \right] \bigvee_a^x(u) \\
 & \quad + \sup_{t \in [x, b]} \left[ (t-x) \left| \int_0^1 f'[\lambda t + (1-\lambda)x] d\lambda \right| \right] \bigvee_x^b(u) \\
 & \leq \sup_{t \in [a, x]} \left[ (x-t) \int_0^1 |f'[\lambda t + (1-\lambda)x]| d\lambda \right] \bigvee_a^x(u) \\
 & \quad + \sup_{t \in [x, b]} \left[ (t-x) \int_0^1 |f'[\lambda t + (1-\lambda)x]| d\lambda \right] \bigvee_x^b(u) \\
 & \leq \sup_{t \in [a, x]} \left[ (x-t) \frac{|f'(t)| + |f'(x)|}{2} \right] \bigvee_a^x(u) + \sup_{t \in [x, b]} \left[ (t-x) \frac{|f'(t)| + |f'(x)|}{2} \right] \bigvee_x^b(u) \\
 & \leq \frac{1}{2} \left[ \sup_{t \in [a, x]} \{ (x-t) |f'(t)| \} \cdot \bigvee_a^x(u) + \sup_{t \in [x, b]} \{ (t-x) |f'(t)| \} \cdot \bigvee_x^b(u) \right] \\
 & \quad + \frac{1}{2} |f'(x)| \left[ (x-a) \bigvee_a^x(u) + (b-x) \bigvee_x^b(u) \right],
 \end{aligned}$$

which proves the first inequality in (3.1).

The second inequality in (3.1) is obvious using properties of sup and the theorem is completely proved. ■

The mid-point inequality is of interest in applications and provides a much simpler inequality:

**Corollary 2.** If  $f$  and  $u$  are as above and  $|f'|$  is convex on  $[a, \frac{a+b}{2}]$  and  $[\frac{a+b}{2}, b]$ , then

$$\begin{aligned}
 & \left| [u(b) - u(a)] f\left(\frac{a+b}{2}\right) - \int_a^b f(t) du(t) \right| \\
 & \leq \frac{1}{4} (b-a) \left[ \|f'\|_{[a, \frac{a+b}{2}], \infty} \bigvee_a^{\frac{a+b}{2}}(u) + \|f'\|_{[\frac{a+b}{2}, b], \infty} \bigvee_{\frac{a+b}{2}}^b(u) \right]
 \end{aligned}$$

$$\begin{aligned}
& + \frac{1}{4} (b-a) \left| f' \left( \frac{a+b}{2} \right) \right| \bigvee_a^b (u) \\
& \leq \frac{1}{4} (b-a) \bigvee_a^b (u) \left[ \|f'\|_{[a,b],\infty} + \left| f' \left( \frac{a+b}{2} \right) \right| \right].
\end{aligned} \tag{3.2}$$

**Remark 4.** If we denote, from the second inequality in (3.1),

$$L_1(x) := \frac{1}{2} \left[ (x-a) \|f'\|_{[a,x],\infty} \bigvee_a^x (u) + (b-x) \|f'\|_{[x,b],\infty} \bigvee_x^b (u) \right]$$

and

$$L_2(x) := \frac{1}{2} |f'(x)| \left[ (x-a) \bigvee_a^x (u) + (b-x) \bigvee_x^b (u) \right]$$

for  $x \in [a, b]$ , then we can point out various upper bounds for the functions  $L_1$  and  $L_2$  on  $[a, b]$ .

For instance, we have

$$L_1(x) \leq \frac{1}{2} \|f'\|_{[a,b],\infty} \left[ (x-a) \bigvee_a^x (u) + (b-x) \bigvee_x^b (u) \right]$$

and by (3.1) we can state the following inequality of interest:

$$\begin{aligned}
& \left| [u(b) - u(a)] f(x) - \int_a^b f(t) \, du(t) \right| \leq \frac{1}{2} [\|f'\|_{[a,b],\infty} + |f'(x)|] \left[ (x-a) \bigvee_a^x (u) + (b-x) \bigvee_x^b (u) \right] \\
& \leq \frac{1}{2} [\|f'\|_{[a,b],\infty} + |f'(x)|] \times \left\{ \left[ \frac{1}{2} (b-a) + \left| x - \frac{a+b}{2} \right| \right] \bigvee_a^b (u) \right. \\
& \quad \left. \left[ \frac{1}{2} \bigvee_a^b (u) + \frac{1}{2} \left| \bigvee_a^x (u) - \bigvee_x^b (u) \right| \right] (b-a) \right\}
\end{aligned} \tag{3.3}$$

for each  $x \in [a, b]$ .

**Remark 5.** A similar result to (3.3) can be stated for the generalised trapezoid rule, out of which we would like to note the following one that is of particular interest:

$$\begin{aligned}
& \left| \int_a^b u(t) \, df(t) - u(b) [f(b) - f(x)] - u(a) [f(x) - f(a)] \right| \\
& \leq \frac{1}{2} [\|f'\|_{[a,b],\infty} + |f'(x)|] \left[ (x-a) \bigvee_a^x (u) + (b-x) \bigvee_x^b (u) \right] \\
& \leq \frac{1}{2} [\|f'\|_{[a,b],\infty} + |f'(x)|] \times \left\{ \left[ \frac{1}{2} (b-a) + \left| x - \frac{a+b}{2} \right| \right] \bigvee_a^b (u) \right. \\
& \quad \left. \left[ \frac{1}{2} \bigvee_a^b (u) + \frac{1}{2} \left| \bigvee_a^x (u) - \bigvee_x^b (u) \right| \right] (b-a) \right\}
\end{aligned} \tag{3.4}$$

for each  $x \in [a, b]$ .

As in Corollary 2, the case  $x = \frac{a+b}{2}$  in (3.4) provides the simple result

$$\left| \int_a^b u(t) \, df(t) - u(b) \left[ f(b) - f\left(\frac{a+b}{2}\right) \right] - u(a) \left[ f\left(\frac{a+b}{2}\right) - f(a) \right] \right|$$



$$\begin{aligned}
&\leq \frac{1}{4} (b-a) \left[ \|f'\|_{[a, \frac{a+b}{2}], \infty} \bigvee_{\frac{a+b}{2}}^{\frac{a+b}{2}}(u) + \|f'\|_{[\frac{a+b}{2}, b], \infty} \bigvee_{\frac{a+b}{2}}^b(u) \right] + \frac{1}{4} (b-a) \left| f' \left( \frac{a+b}{2} \right) \right| \bigvee_a^b(u) \\
&\leq \frac{1}{4} (b-a) \bigvee_a^b(u) \left[ \|f'\|_{[a, b], \infty} + \left| f' \left( \frac{a+b}{2} \right) \right| \right]. \tag{3.5}
\end{aligned}$$

**Remark 6.** Similar inequalities may be stated if one assumes either that  $|f'|$  is quasi-convex or that  $|f'|$  is log-convex on  $[a, x]$  and  $[x, b]$ . The details are left to the interested readers.

### Acknowledgements

The authors would like to thank the anonymous referee for valuable suggestions that have been implemented in the final version of the manuscript. The second author's research is supported in part by the Research Grants Council of the Hong Kong SAR (Project No. HKU7017/05P).

### References

- [1] A. Ostrowski, Über die absolutabweichung einer differentienbaren Funktionen von ihren integralmittelwert, *Comment. Math. Helv* 10 (1938) 226–227.
- [2] S.S. Dragomir, S. Wang, A new inequality of Ostrowski's type in  $L_1$ -norm and applications to some special means and to some numerical quadrature rules, *Tamkang J. Math.* 28 (1997) 239–244.
- [3] S.S. Dragomir, S. Wang, A new inequality of Ostrowski's type in  $L_p$ -norm and applications to some special means and to some numerical quadrature rules, *Indian J. Math.* 40 (3) (1998) 245–304.
- [4] S.S. Dragomir, S. Wang, Applications of Ostrowski's inequality to the estimation of error bounds for some special means and some numerical quadrature rules, *Appl. Math. Lett.* 11 (1998) 105–109.
- [5] A.M. Fink, Bounds on the deviation of a function from its averages, *Czechoslovak Math. J.* 42(117) (2) (1992) 298–310.
- [6] S.S. Dragomir, The Ostrowski's integral inequality for Lipschitzian mappings and applications, *Comput. Math. Appl.* 38 (1999) 33–37.
- [7] S.S. Dragomir, P. Cerone, J. Roumeliotis, S. Wang, A weighted version of Ostrowski inequality for mappings of Hölder type and applications in numerical analysis, *Bull. Math. Soc. Sci. Math. Roumanie* 42(90) (4) (1999) 301–314.
- [8] S.S. Dragomir, On the Ostrowski's inequality for mappings of bounded variation and applications, *Math. Inequal. Appl.* 4 (1) (2001) 33–40.
- [9] S.S. Dragomir, Ostrowski's inequality for monotonous mappings and applications, *J. KSIAM* 3 (1) (1999) 127–135.
- [10] S.S. Dragomir, An Ostrowski like inequality for convex functions and applications, *Rev. Mat. Complut.* 16 (2) (2003) 373–382.
- [11] S.S. Dragomir, On the Ostrowski's inequality for Riemann–Stieltjes integral, *Korean J. Appl. Math.* 7 (2000) 477–485.
- [12] S.S. Dragomir, On the Ostrowski inequality for Riemann–Stieltjes integral  $\int_a^b f(t) du(t)$  where  $f$  is of Hölder type and  $u$  is of bounded variation and applications, *J. KSIAM* 5 (1) (2001) 35–45.
- [13] G.A. Anastassiou, Univariate Ostrowski inequalities, *Monatsh. Math.* 135 (3) (2002) 175–189. Revisited.
- [14] P. Kumar, The Ostrowski type moments integral inequalities and moment bounds for continuous random variables, *Comput. Math. Appl.* 49 (11–12) (2005) 1929–1940.
- [15] Z. Liu, Refinement of an inequality of Grüss type for Riemann–Stieltjes integral, *Soochow J. Math.* 30 (4) (2004) 483–489.
- [16] P. Cerone, S.S. Dragomir, New bounds for the three-point rule involving the Riemann–Stieltjes integrals, in: C. Gulati, et al. (Eds.), *Advances in Statistics Combinatorics and Related Areas*, World Science Publishing, 2002, pp. 53–62.
- [17] S.S. Dragomir, Ostrowski type inequalities for isotonic linear functionals, *J. Inequal. Pure Appl. Math.* 3 (5) (2002). Online: <http://jipam.vu.edu.au/article.php?sid=220>.